

Roll No.-----

Paper Code		
3	7	0
(To be filled in the OMR Sheet)		

प्रश्नपुस्तिका क्रमांक
Question Booklet No.

O.M.R. Serial No.

--	--	--	--	--	--	--	--

प्रश्नपुस्तिका सीरीज
Question Booklet Series
D

BCA (Second Semester) Examination, July-2022

BCA-205(N)

Mathematics-II
(B.P.)

Time : 1:30 Hours

Maximum Marks-100

जब तक कहा न जाय, इस प्रश्नपुस्तिका को न खोलें

- निर्देश : —**
1. परीक्षार्थी अपने अनुक्रमांक, विषय एवं प्रश्नपुस्तिका की सीरीज का विवरण यथास्थान सही- सही भरें, अन्यथा मूल्यांकन में किसी भी प्रकार की विसंगति की दशा में उसकी जिम्मेदारी स्वयं परीक्षार्थी की होगी।
 2. इस प्रश्नपुस्तिका में 100 प्रश्न हैं, जिनमें से केवल 75 प्रश्नों के उत्तर परीक्षार्थियों द्वारा दिये जाने हैं। प्रत्येक प्रश्न के चार वैकल्पिक उत्तर प्रश्न के नीचे दिये गये हैं। इन चारों में से केवल एक ही उत्तर सही है। जिस उत्तर को आप सही या सबसे उचित समझते हैं, अपने उत्तर पत्रक (O.M.R. ANSWER SHEET) में उसके अक्षर वाले वृत्त को काले या नीले बाल प्वाइंट पेन से पूरा भर दें। यदि किसी परीक्षार्थी द्वारा किसी प्रश्न का एक से अधिक उत्तर दिया जाता है, तो उसे गलत उत्तर माना जायेगा।
 3. प्रत्येक प्रश्न के अंक समान हैं। आप के जितने उत्तर सही होंगे, उन्हीं के अनुसार अंक प्रदान किये जायेंगे।
 4. सभी उत्तर केवल ओ०एम०आर० उत्तर पत्रक (O.M.R. ANSWER SHEET) पर ही दिये जाने हैं। उत्तर पत्रक में निर्धारित स्थान के अलावा अन्यत्र कहीं पर दिया गया उत्तर मान्य नहीं होगा।
 5. ओ०एम०आर० उत्तर पत्रक (O.M.R. ANSWER SHEET) पर कुछ भी लिखने से पूर्व उसमें दिये गये सभी अनुदेशों को सावधानीपूर्वक पढ़ लिया जाय।
 6. परीक्षा समाप्ति के उपरान्त परीक्षार्थी कक्ष निरीक्षक को अपनी ओ०एम०आर० शीट उपलब्ध कराने के बाद ही परीक्षा कक्ष से प्रस्थान करें।
 7. निगेटिव मार्किंग नहीं है।

महत्वपूर्ण : — प्रश्नपुस्तिका खोलने पर प्रथमतः जाँच कर देख लें कि प्रश्नपुस्तिका के सभी पृष्ठ भलीभाँति छपे हुए हैं। यदि प्रश्नपुस्तिका में कोई कमी हो, तो कक्ष निरीक्षक को दिखाकर उसी सीरीज की दूसरी प्रश्नपुस्तिका प्राप्त कर लें।

K-370

1. Triple integral is used to calculate :
 - (A) Area
 - (B) Volume
 - (C) (A) only
 - (D) NOT
2. Changing the order of integration the integral $\int_{y=2}^3 \int_{x=0}^2 f(x, y) dx dy$ is equal to :
 - (A) $\int_{x=0}^2 \int_{y=2}^3 f(x, y) dy dx$
 - (B) $\int_{x=0}^3 \int_{y=2}^2 f(x, y) dy dx$
 - (C) (A) and (B) both
 - (D) NOT
3. Using volume integral, which of the following quantity can be calculated :
 - (A) Area of circle
 - (B) Area of square
 - (C) Volume of cube
 - (D) NOT
4. Double integral $\int_{x_1}^{x_2} \int_{y_1}^{y_2} dy dx$ represents :
 - (A) Volume
 - (B) Area and volume
 - (C) Area
 - (D) NOT
5. $\int_a^b \int_c^d dx dy$ is :
 - (A) $-\int_c^d \int_a^b dy dx$
 - (B) $\int_c^d \int_a^b dy dx$
 - (C) $abcd$
 - (D) $bc - ad$

6. Distance of plane $3x + 4y - 5z - 7 = 0$ from origin is :
- (A) $\frac{7}{\sqrt{40}}$
 (B) $\frac{6}{\sqrt{34}}$
 (C) $\frac{8}{\sqrt{50}}$
 (D) $\frac{7}{\sqrt{50}}$
7. If l, m, n represents the direction cosines then $l^2 + m^2 + n^2$ is :
- (A) 0
 (B) 3
 (C) -1
 (D) 1
8. If $u(x, y) = xy + yz + zx$, then $\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} + \frac{\partial f}{\partial z}$ is :
- (A) $y + x$
 (B) $y + z$
 (C) $2(x + y + z)$
 (D) $x + y + z$
9. Equation of sphere in standard form is :
- (A) $x^2 + y^2 + z^2 = r^2$
 (B) $x^2 + y^2 + 2gx + 2fy + c = 0$
 (C) $xyz = r^2$
 (D) $(x - h)^2 + (y - k)^2 = r^2$
10. If the lines $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$ and $\frac{x-x_2}{p} = \frac{y-y_2}{q} = \frac{z-z_2}{r}$ lie on a plane then :
- (A) $\begin{vmatrix} x_1 & y_1 & z_1 \\ a & b & c \\ p & q & r \end{vmatrix} = 0$
 (B) $\begin{vmatrix} x_1 - x_2 & y_1 - y_2 & z_1 - z_2 \\ a & b & c \\ p & q & r \end{vmatrix} = 0$
 (C) $\begin{vmatrix} x_1 - x_2 & y_1 - y_2 & z_1 - z_2 \\ a & b & c \\ p & q & r \end{vmatrix} \neq 0$
 (D) NOT

11. $A \times B$ is defined as :
- (A) $\{(a, b) ; a, b \in A\}$
 - (B) $\{(a, b) ; a, b \in B\}$
 - (C) $\{(a, b) ; a \in A, b \in B\}$
 - (D) $\{(a, b) ; a \in B, b \in A\}$
12. Set of integers z with relation ' $>$ ' is a :
- (A) POSET
 - (B) Not POSET but Reflexive
 - (C) Reflexive
 - (D) Not a POSET
13. If $A = \{x; x^2 \leq 16, x \in \mathbb{Z}\}$ then A is :
- (A) $\{0, 1, 2, 3, 4\}$
 - (B) $\{-4, -3, -2, -1, 0, 1, 2, 3, 4\}$
 - (C) $\{x; -4 \leq x \leq 4\}$
 - (D) NOT
14. If $P(S)$ is power set of S , then $(P(S), \subseteq)$ is :
- (A) Never a lattice
 - (B) Always a lattice
 - (C) $|P(S)| = |S|$
 - (D) $|P(S)| = 2$
15. Relation ' $<$ ' on $\mathbb{Z}^+$ is :
- (A) Partial order relation
 - (B) Not a partial order relation
 - (C) Reflexive
 - (D) NOT
16. Set $A = \{x; 3x - 2 = 3, x \in \mathbb{Z}\}$ is :
- (A) $\left\{\frac{5}{2}\right\}$
 - (B) $\frac{5}{3}$
 - (C) ϕ
 - (D) $\{\phi\}$

17. If $A = \{a, b\}, B = \{b, a\}$ then :
- (A) $A \neq B$
 - (B) $A = B$
 - (C) $A \cap B = \phi$
 - (D) NOT
18. If $f(x, y) = x^3 - 3xy^2, g(x, y) = 3x^2y - y^3$ then :
- (A) $\frac{\partial f}{\partial x} = \frac{\partial g}{\partial x}$
 - (B) $\frac{\partial f}{\partial x} = \frac{-\partial g}{\partial y}$
 - (C) $\frac{\partial f}{\partial y} = \frac{\partial g}{\partial x}$
 - (D) $\frac{\partial f}{\partial y} = \frac{-\partial g}{\partial x}$
19. $n(A \cup B)$ is equal to :
- (A) $n(A) + n(B) - n(A \cap B)$
 - (B) $n(A) + n(B)$
 - (C) $n(A) - n(B)$
 - (D) $n(A) \cup n(B)$
20. If $n(A) = 20, n(B) = 30, n(A \cup B) = 100$ then $n(A \cap B)$ is :
- (A) 50
 - (B) 40
 - (C) 20
 - (D) 30
21. $A = \{1, 2, 3\}, B = \{4, 5, 6\}$ then $A - B$ is :
- (A) $\{1, 2, 3\}$
 - (B) $\{4, 5, 6\}$
 - (C) ϕ
 - (D) B

22. If $f(x, y) = \frac{x^3+y^3}{\sqrt{x}+\sqrt{y}}$ is homogeneous function of degree n then n is :
- (A) $\frac{3}{2}$
 - (B) 3
 - (C) $\frac{5}{2}$
 - (D) $\frac{1}{2}$
23. If $u = e^{xyz}$ then $\frac{\partial f}{\partial x}$ is :
- (A) yzu
 - (B) $yz e^{yz}$
 - (C) xyz
 - (D) xyu
24. Set of circles passing through origin is a :
- (A) Finite set
 - (B) Infinite set
 - (C) Both finite and infinite set
 - (D) NOT
25. $(A \cup B)'$ is equal to :
- (A) $A' \cup B'$
 - (B) $A' \cup \overline{B}$
 - (C) $A' \cap B'$
 - (D) $(A \cap B)'$
26. If $A = \phi$ then power set of A is :
- (A) $\{\phi\}$
 - (B) $\{\phi, A\}$
 - (C) ϕ
 - (D) NOT

27. Number of elements in power set of set $A = \{1, 2, 3\}$ is :
- (A) 3
(B) 8
(C) 9
(D) 6
28. If u is homogeneous function of x, y with degree n then :
- (A) $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = nu$
(B) $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = u$
(C) $x \frac{\partial u}{\partial x} - y \frac{\partial u}{\partial y} = nu$
(D) NOT
29. If $z = \tan^{-1} \left(\frac{y}{x} \right)$, then the value of $\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2}$ is equal to :
- (A) $\frac{-y}{x^2+y^2}$
(B) $\frac{x}{x^2+y^2}$
(C) $\frac{2xy}{x^2+y^2}$
(D) 0
30. If $y = \log \sin x + z$ then $\frac{\partial y}{\partial x}$ is :
- (A) $\cot x$
(B) $\frac{1}{\sin x}$
(C) $\cos x$
(D) z
31. If $f(x, y, z) = x^2 + xyz + z$ then $f_x(1, 1, 1)$ is :
- (A) 0
(B) 3
(C) 1
(D) -1

32. If a line has direction ratio's (1, 2, 3) then its direction cosines are :

(A) $\left(\frac{1}{\sqrt{14}}, \frac{-2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

(B) $\left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

(C) $\left(\frac{-1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

(D) $\left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{-3}{\sqrt{14}}\right)$

33. Sum of direction cosines of z axis is :

(A) 0

(B) $\frac{1}{3}$

(C) 1

(D) 3

34. Volume of a cube with side a is :

(A) $\int_0^a \int_0^a \int_0^a dx dy dz$

(B) a^2

(C) $\int_0^a \int_0^a dx dy$

(D) $\frac{a^3}{8}$

35. To find the volume, which of the following integral can be used :

(A) Single

(B) Double

(C) Triple

(D) Double and triple

36. $\int_0^1 \int_0^x dx dy$ is :

(A) 1

(B) $\frac{1}{2}$

(C) 2

(D) 3

37. Changing the order of integration in $\int_0^a \int_0^y f(x, y) dx dy$, the new integral is :
- (A) $\int_0^a \int_x^a f(x, y) dy dx$
- (B) $\int_x^a \int_0^a f(x, y) dy dx$
- (C) $\int_0^a \int_x^y f(x, y) dy dx$
- (D) $\int_0^a \int_{a-x}^a f(x, y) dy dx$
38. If $u = \log \left[\frac{x^2+y^2}{x+y} \right]$, then $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$ is :
- (A) 0
- (B) 1
- (C) u
- (D) eu
39. If the lines $3x + 2ky - 2 = 0$ and $2x + 5y + 1 = 0$ are parallel then the value of k is :
- (A) $\frac{-5}{4}$
- (B) $\frac{3}{2}$
- (C) $\frac{15}{4}$
- (D) $\frac{2}{5}$
40. Which of the following is true ?
- (A) $\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial y^2}$
- (B) $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$
- (C) $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$
- (D) NOT

41. If $f(x, y) = \sqrt{x} + \sqrt{y}$ then degree of Homogeneous function $f(x, y)$ is :
- (A) $\frac{1}{2}$
 - (B) 2
 - (C) $\frac{1}{4}$
 - (D) NOT
42. $\int_0^\pi \int_0^\pi d\theta d\phi$ is :
- (A) 1
 - (B) 0
 - (C) $\frac{\pi}{2}$
 - (D) π^2
43. Curve $y^2 = 4x$ is a :
- (A) Parabola
 - (B) Hyperbola
 - (C) Straight line
 - (D) Ellipse
44. $\int_0^1 \int_0^1 \int_0^1 dx dy dz$ is :
- (A) 1
 - (B) 2
 - (C) 0
 - (D) 4
45. A line makes an angle α, β, γ , with x, y, z , axis respectively then $\sin^2\alpha + \sin^2\beta + \sin^2\gamma$ is :
- (A) 1
 - (B) 0
 - (C) 2
 - (D) NOT

46. Direction ratios of normal to the plane $2x - y + 2z + 1 = 0$ are :
- (A) $(2, 1, 2)$
 - (B) $(2, -1, 2)$
 - (C) $(2, 1, -2)$
 - (D) NOT
47. Acute angle between the lines $x - 2 = 0$ and $\sqrt{3}x - y - 2 = 0$ is :
- (A) 0°
 - (B) 30°
 - (C) 45°
 - (D) 60°
48. Equation of sphere with center at $(-3, 2, 4)$ and radius 6 units is :
- (A) $x^2 + y^2 + z^2 + 6x + 4y + 8z = 7$
 - (B) $x^2 + y^2 + z^2 + 6x - 4y - 8z = 7$
 - (C) $x^2 + y^2 + z^2 - 6x + 4y - 8z = 7$
 - (D) None
49. If $u = \frac{x^4 + y^4}{x + y}$, then $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$ is :
- (A) $4u$
 - (B) 4
 - (C) $3u$
 - (D) NOT
50. If $f(x, y) = x^2 + y^2$, then $\frac{\partial f}{\partial x}$ is :
- (A) $2x + y^2$
 - (B) $2x$
 - (C) $2x + 2y$
 - (D) NOT

51. Range of function $f(x) = \tan x$ is :
- (A) $(-1, 1)$
 - (B) $\left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$
 - (C) $[-\pi, \pi]$
 - (D) $(-\infty, \infty)$
52. Equation of a plane passing through $(1, 2, 1), (2, -1, -4), (1, 0, -1)$ is :
- (A) $2x - y + z = 1$
 - (B) $x + y + z = 1$
 - (C) $2x + y + z = 1$
 - (D) NOT
53. Function $f: N \rightarrow N$ such that $f(x) = x + 1$ is :
- (A) One-one only
 - (B) Onto only
 - (C) One-one onto
 - (D) NOT
54. Range of $f(x) = \cos x$ is :
- (A) $(-1, 1)$
 - (B) $[-2, 2]$
 - (C) $[-\pi, \pi]$
 - (D) $[-1, 1]$
55. Stationary point is a point where $f(x, y)$ have ?
- (A) $\frac{\partial f}{\partial x} = 0$
 - (B) $\frac{\partial f}{\partial y} = 0$
 - (C) $\frac{\partial f}{\partial x} = 0$ and $\frac{\partial f}{\partial y} = 0$
 - (D) $\frac{\partial f}{\partial x} < 0$ and $\frac{\partial f}{\partial y} > 0$
56. Saddle point is a point where function has :
- (A) Maximum value
 - (B) Minimum value
 - (C) Zero value
 - (D) Neither maximum nor minimum value


57. The value of $\int_0^2 \int_1^y xy \, dx \, dy$ is :

- (A) 0
- (B) -1
- (C) 2
- (D) 1

58. Let $D_{30} = \{1, 2, 3, 5, 6, 10, 15, 30\}$ and relation 'a divides b' be a partial ordering on D_{30} . The lub of 10 and 15 respectively is :


- (A) 30
- (B) 15
- (C) 10
- (D) 16

59. Which element is minimal in the following diagram :



- (A) d
- (B) b
- (C) c
- (D) a

60. The graph given below is an example of :



- (A) Non lattice
- (B) Lattice
- (C) Semi lattice
- (D) NOT

61. Function $f(x, y)$ has maximum value at (a, b) if :
- (A) $rt - s^2 > 0$ and $r < 0$
 - (B) $rt - s^2 > 0$ and $r > 0$
 - (C) $rt - s^2 < 0$ and $r < 0$
 - (D) $rt - s^2 < 0$ and $r > 0$
62. If $u = \log(x + y + 1)$, then $\frac{\partial u}{\partial x}$ at $(1, 2)$:
- (A) $\frac{1}{3}$
 - (B) 4
 - (C) $\frac{1}{2}$
 - (D) $\frac{1}{4}$
63. Let $X = \{2, 3, 6, 12, 24\}$ and ' $\leq$ ' be the partial order relation $x \leq y$ if x divides y . Number of edges in the Hasse diagram of $(X, \leq)$ is :
- (A) 3
 - (B) 4
 - (C) 9
 - (D) NOT
64. If $f(x) = \frac{2x+5}{x^2+x+5}$, then $f(f(-1))$ is equal to :
- (A) $\frac{155}{149}$
 - (B) $\frac{149}{155}$
 - (C) 0
 - (D) NOT
65. If the function $f: Q \rightarrow Q$ is defined by the relation $f(x) = x - \frac{1}{2}$, $x \in Q$ then f is :
- (A) Many one onto mapping
 - (B) One-one into mapping
 - (C) One-one onto mapping
 - (D) Many one onto mapping

66. $A = \{1, 2, 3, 4\}$, $B = \{2, 3, 4, 5\}$, $C = \{4, 5, 6, 7\}$ then $(A \cap B) \cap C$ then :
- (A) $\{4\}$
 - (B) $\{2, 3, 4\}$
 - (C) $\{3, 4\}$
 - (D) ϕ
67. If R is an equivalence relation on set B then R^{-1} is :
- (A) Reflexive only
 - (B) Symmetric only
 - (C) Equivalence
 - (D) NOT
68. Let R and T be two equivalence relations on a set A then :
- (A) $R \cup S$ is an equivalence relation
 - (B) $R \cap S$ is an equivalence relation
 - (C) $R - S$ is an equivalence relation
 - (D) NOT
69. Let R be a reflexive relation on a set A and I be the identity relation on A , then :
- (A) $R \subset I$
 - (B) $I \subset R$
 - (C) $R = I$
 - (D) NOT
70. A relation $f(x) = y$ s.t. :
- (A) $y^2 = 4ax$ is a function
 - (B) $y^2 = -4ax$ is a function
 - (C) $y = x$ is a function
 - (D) NOT
71. If $f(x) = \frac{x+2}{x-3}$, $x \neq 3$ then $f^{-1}(x)$ is equal to :
- (A) $\frac{x-1}{3x+2}$
 - (B) $\frac{3x+2}{x-1}$
 - (C) $\frac{x-3}{x+2}$
 - (D) $\frac{x-2}{x+3}$

72. Range of function $f: R \rightarrow R$ st. $f(1) = 2, f(2) = 3, f(3) = 4$ is :
- (A) $\{1, 2, 3\}$
 (B) R
 (C) $\{2, 3, 4\}$
 (D) $\{(1, 2), (2, 3), (3, 4)\}$
73. The number of elements in the power set of a set having $n - 2$ elements is :
- (A) $2^n - 2$
 (B) 2^n
 (C) 2^{n-2}
 (D) 2^{n-1}
74. Relation $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$ on the set $A = \{1, 2, 3\}$ is :
- (A) Reflexive but not symmetric
 (B) Reflexive but not transitive
 (C) Symmetric and transitive
 (D) NOT
75. If $R \subset A \times B$ and $S \subset B \times C$ be two relations, then $(SoR)^{-1}$ equals to :
- (A) $S^{-1} \circ R^{-1}$
 (B) $R^{-1} \circ S^{-1}$
 (C) SoR
 (D) RoS
76. Let $f: z \rightarrow z$ (z is set of integers) be defined by $f(x) = x^2 + x - 2$ then $f(f(-2))$ is :
- (A) -2
 (B) -1
 (C) 1
 (D) 3
77. Let $A = \{-2, -1, 0\}$ and $f(x) = 2x - 3$ then the range of f is :
- (A) $\{7, -5, -3\}$
 (B) $(-7, 5, -3)$
 (C) $\{-7, -5, -3\}$
 (D) NOT

78. If $f(x) = x^2$ and $g(x) = \sin x$ then the value of $g \circ f(x)$ is :
- (A) $\cos x$
 - (B) $\sin x^2$
 - (C) $-\sin x^2$
 - (D) $\sin x$
79. Which of the following is a partial order as well as equivalence relation?
- (A) Equal to ($=$)
 - (B) Less than ($<$)
 - (C) Greater than ($>$)
 - (D) NOT
80. Which laws are satisfied for a lattice?
- (A) Associative law
 - (B) Commutative law
 - (C) Absorption law
 - (D) All above
81. A POSET in which every pair of element has both a least upper bound and greatest lower bound is :
- (A) Sublattice
 - (B) Lattice
 - (C) Not a lattice
 - (D) NOT
82. $a \wedge (b \wedge c) = (a \wedge b) \wedge c$ is :
- (A) Distributive law
 - (B) Associative law
 - (C) commutative law
 - (D) NOT

83. Maximum value of xe^{-x} is :
- (A) e
 - (B) $\frac{1}{e}$
 - (C) $-e$
 - (D) $-\frac{1}{e}$
84. If $f(x, y) = \frac{x+y}{y}$ then $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y}$ is :
- (A) 0
 - (B) 1
 - (C) 2
 - (D) 3
85. If $u = ax^2 + 2hxy + by^2$ then using Euler's theorem, $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$ is :
- (A) u
 - (B) $2u$
 - (C) $3u$
 - (D) $n(n-1)$
86. Domain of the function $f(x) = \sqrt{4-x^2}$ is all real x s.t. :
- (A) $x < 2$
 - (B) -2
 - (C) $-2 \leq x \leq 2$
 - (D) $x > -2$
87. A function $f: R \rightarrow R$ s.t. $f(x) = x^2$ is :
- (A) One one, onto
 - (B) One one, into
 - (C) Many one into
 - (D) Does not exist

88. Domain of $\sin^{-1}(4x)$ is :
- (A) $[0, 1]$
 - (B) $\left[\frac{-1}{4}, \frac{1}{4}\right]$
 - (C) $[-3, 3]$
 - (D) NOT
89. The relation ' $<$ ' in the set of natural numbers is :
- (A) Only symmetric
 - (B) Only transitive
 - (C) Only reflexive
 - (D) Equivalence relation
90. Let $f: R \rightarrow R$ be defined by $f(x) = 3x - 4$, then $f^{-1}(x)$ is :
- (A) $\frac{x+4}{3}$
 - (B) $\frac{x}{3} - 4$
 - (C) $3x + 4$
 - (D) $x + 3$
91. Radius of the sphere $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ is :
- (A) $\sqrt{u^2 + v^2 + w^2}$
 - (B) $\sqrt{u^2 + v^2 + w^2 - d}$
 - (C) $\sqrt{u^2 + v^2 + w^2 + d}$
 - (D) NOT
92. Center of sphere $x^2 + y^2 + z^2 + 2x - 4y - 6z + 5 = 0$ is :
- (A) $(1, 2, 3)$
 - (B) $(-1, -2, 3)$
 - (C) $(-1, 2, 3)$
 - (D) $(1, 2, -3)$

93. If $A = \{a, b\}$ then power set $P(A)$ is :
- (A) $\{\phi, \{a\}, \{b\}, \{a, b\}\}$
 - (B) $\{\{a\}, \{b\}\}$
 - (C) $\{\{a\}, \{b\}, \{a, b\}\}$
 - (D) $\{\phi, a, b, A\}$
94. If $A = \{x: x \text{ is a factor of } 10\}$ $B = \{2, 5\}$ then :
- (A) $A \subset B$
 - (B) $A = B$
 - (C) $B \subset A$
 - (D) $A \cap B = \phi$
95. If $n(A) = 40$, $n(B) = 40$, $n(A \cap B) = 20$, $n(A \cup B) = 60$ then $n(B \setminus A)$ is :
- (A) 60
 - (B) 40
 - (C) 30
 - (D) 20
96. If $A = \{1\}$, $B = \{2\}$ then $A \times B$ is :
- (A) $\{(1, 2)\}$
 - (B) $\{1, 2\}$
 - (C) $\{1\}$
 - (D) $\{2\}$
97. If $A_i = \{0, i\}, i \in \mathbb{Z}$ then $A_1 \cup A_2$ is equal to :
- (A) A_1
 - (B) A_2
 - (C) $A_1 \cap A_2$
 - (D) A_3

98. Which of the following represent the equation of plane?
- (A) $ax^2 + by = d$
 - (B) $ax + by + cz = xy$
 - (C) $ax + by + cz = d$
 - (D) NOT
99. Mid points of the line segment joining the points $(2, 3, 4)$ and $(8, -3, 8)$ are :
- (A) $(10, 0, 12)$
 - (B) $(5, 6, 0)$
 - (C) $(6, 5, 0)$
 - (D) $(5, 0, 6)$
100. If the plane passes through three collinear points $(x_1, y_1, z_1), (x_2, y_2, z_2), (x_3, y_3, z_3)$ then which of the following true ?
- (A) $x_1y_1z_1 + x_2y_2z_2 + x_3y_3z_3 = 0$
 - (B) $\begin{vmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix} = 0$
 - (C) $x_1 + y_1 + z_1 = 0$
 - (D) NOT

Rough Work / रफ कार्य

DO NOT OPEN THE QUESTION BOOKLET UNTIL ASKED TO DO SO

1. Examinee should enter his / her roll number, subject and Question Booklet Series correctly in the O.M.R. sheet, the examinee will be responsible for the error he / she has made.
 2. **This Question Booklet contains 100 questions, out of which only 75 Question are to be Answered by the examinee. Every question has 4 options and only one of them is correct. The answer which seems correct to you, darken that option number in your Answer Booklet (O.M.R ANSWER SHEET) completely with black or blue ball point pen. If any examinee will mark more than one answer of a particular question, then the answer will be marked as wrong.**
 3. Every question has same marks. Every question you attempt correctly, marks will be given according to that.
 4. Every answer should be marked only on Answer Booklet (O.M.R ANSWER SHEET). Answer marked anywhere else other than the determined place will not be considered valid.
 5. Please read all the instructions carefully before attempting anything on Answer Booklet **(O.M.R ANSWER SHEET)**.
 6. After completion of examination, please hand over the **O.M.R. SHEET** to the Examiner before leaving the examination room.
 7. There is no negative marking.
- Note:** On opening the question booklet, first check that all the pages of the question booklet are printed properly in case there is an issue please ask the examiner to change the booklet of same series and get another one.