

Roll No.-----

Paper Code		
3	7	6
(To be filled in the OMR Sheet)		

प्रश्नपुस्तिका क्रमांक
Question Booklet No.

O.M.R. Serial No.

--	--	--	--	--	--	--	--

प्रश्नपुस्तिका सीरीज
Question Booklet Series
B

BCA (Fourth Semester) Examination, July-2022

BCA-405(N)

Mathematics-III

Time : 1:30 Hours

Maximum Marks-100

जब तक कहा न जाय, इस प्रश्नपुस्तिका को न खोलें

- K-376**
- निर्देश : —
- परीक्षार्थी अपने अनुक्रमांक, विषय एवं प्रश्नपुस्तिका की सीरीज का विवरण यथास्थान सही- सही भरें, अन्यथा मूल्यांकन में किसी भी प्रकार की विसंगति की दशा में उसकी जिम्मेदारी स्वयं परीक्षार्थी की होगी।
 - इस प्रश्नपुस्तिका में 100 प्रश्न हैं, जिनमें से केवल 75 प्रश्नों के उत्तर परीक्षार्थियों द्वारा दिये जाने हैं। प्रत्येक प्रश्न के चार वैकल्पिक उत्तर प्रश्न के नीचे दिये गये हैं। इन चारों में से केवल एक ही उत्तर सही है। जिस उत्तर को आप सही या सबसे उचित समझते हैं, अपने उत्तर पत्रक (O.M.R. ANSWER SHEET) में उसके अक्षर वाले वृत्त को काले या नीले बाल प्वाइंट पेन से पूरा भर दें। यदि किसी परीक्षार्थी द्वारा किसी प्रश्न का एक से अधिक उत्तर दिया जाता है, तो उसे गलत उत्तर माना जायेगा।
 - प्रत्येक प्रश्न के अंक समान हैं। आप के जितने उत्तर सही होंगे, उन्हीं के अनुसार अंक प्रदान किये जायेंगे।
 - सभी उत्तर केवल ओ०एम०आर० उत्तर पत्रक (O.M.R. ANSWER SHEET) पर ही दिये जाने हैं। उत्तर पत्रक में निर्धारित स्थान के अलावा अन्यत्र कहीं पर दिया गया उत्तर मान्य नहीं होगा।
 - ओ०एम०आर० उत्तर पत्रक (O.M.R. ANSWER SHEET) पर कुछ भी लिखने से पूर्व उसमें दिये गये सभी अनुदेशों को सावधानीपूर्वक पढ़ लिया जाय।
 - परीक्षा समाप्ति के उपरान्त परीक्षार्थी कक्ष निरीक्षक को अपनी ओ०एम०आर० शीट उपलब्ध कराने के बाद ही परीक्षा कक्ष से प्रस्थान करें।
 - निगेटिव मार्किंग नहीं है।
- महत्वपूर्ण : — प्रश्नपुस्तिका खोलने पर प्रथमतः जाँच कर देख लें कि प्रश्नपुस्तिका के सभी पृष्ठ भलीभाँति छपे हुए हैं। यदि प्रश्नपुस्तिका में कोई कमी हो, तो कक्ष निरीक्षक को दिखाकर उसी सीरीज की दूसरी प्रश्नपुस्तिका प्राप्त कर लें।

1. In sequence $\langle a_n \rangle$, where $a_n = \frac{1}{n}$ least upper bound is :
 - (A) 0
 - (B) 1
 - (C) 2
 - (D) -1
2. Sequence $\langle 1 + \frac{1}{n} \rangle$ is :
 - (A) Bounded
 - (B) Not bounded
 - (C) Bounded above only
 - (D) Bounded below only
3. Sequence $\langle 2 + \frac{7}{n+5} \rangle$ converges to :
 - (A) 7
 - (B) 0
 - (C) 5
 - (D) 2
4. If sequence, $\langle a_n \rangle$ is bounded above there exist a real number k s.t. :
 - (A) $a_n \leq k, \forall n \in N$
 - (B) $a_n \geq k, \forall n \in N$
 - (C) $a_n < -k, \forall n \in N$
 - (D) NOT
5. Series $\sum_{n=1}^{\infty} \frac{1}{n}$ is :
 - (A) Convergent
 - (B) Divergent
 - (C) Oscillatory
 - (D) NOT

6. If series $\sum_{n=1}^{\infty} a_n$ is convergent then :

(A) $\lim_{n \rightarrow \infty} a_n = 0$

(B) $\lim_{n \rightarrow \infty} a_n \neq 0$

(C) $\lim_{n \rightarrow \infty} a_n = 1$

(D) $\lim_{n \rightarrow \infty} a_n =$ does not exist

7. Series $\sum_{n=1}^{\infty} \cos \frac{1}{n}$ is :

(A) Convergent

(B) Divergent

(C) Oscillatory

(D) NOT

8. Series $\sum_{n=0}^{\infty} x^n$ is convergent if :

(A) $|x| = 1$

(B) $|x| > 1$

(C) $|x| < 1$

(D) $x = 1$

9. p – series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is divergent :

(A) $p = 2$

(B) $p = 3$

(C) $p = 4$

(D) $p = 0$

10. $\sum_{n=1}^{\infty} \frac{1}{n^4}$ is :
(A) Divergent series
(B) Oscillatory series
(C) Convergent series
(D) NOT
11. Field $\vec{F} = 2xy\hat{i} + (x^2 + z^2)\hat{j} + 2zy\hat{k}$ is :
(A) Solenoidal
(B) Rotational
(C) Irrotational
(D) NOT
12. Curl of a vector field $\vec{f}(x, y, z) = x^2\hat{i} + 2z\hat{j} - y\hat{k}$ is :
(A) $-3\hat{i}$
(B) $-3\hat{j}$
(C) $-3\hat{k}$
(D) 0
13. If $\text{curl}(\vec{v}) = 0$ then field $\vec{v}$ is :
(A) Solenoidal
(B) Rotational
(C) Irrotational
(D) NOT
14. Particular integral for $(D^2 + 1)y = e^{2x}$ is :
(A) $\frac{1}{4}e^{2x}$
(B) $\frac{1}{5}e^{2x}$
(C) $\frac{1}{2}e^{2x}$
(D) $\frac{1}{6}e^{2x}$

15. Particular Integral for $(D^2 + 2D + 1)y = 0$ is :
- (A) $\frac{1}{2}$
 - (B) 0
 - (C) 2
 - (D) NOT
16. Solution of $(D^2 + 5D + 4)y = 0$ is :
- (A) $c_1 e^x + c_2 e^{-4x}$
 - (B) $c_1 e^{-x} + c_2 e^{4x}$
 - (C) $c_1 e^{-x} + c_2 e^{-4x}$
 - (D) NOT
17. $\frac{dy}{dx} = \frac{ax+by+c}{dx+ey+f}$ is transformed into linear ODE by taking $x = X + h$, $y = Y + K$ if :
- (A) $ae - bd = 0$
 - (B) $ae - bd \neq 0$
 - (C) $ad - be \neq 0$
 - (D) $ad - be = 0$
18. Solution of $\frac{dy}{dx} = \frac{x}{y}$ is :
- (A) $y^2 = x^2 + c$
 - (B) $y^2 = -x^2 + c$
 - (C) $y = x + c$
 - (D) $y = x - c$
19. $y - x \frac{dy}{dx} = x + y \frac{dy}{dx}$ is :
- (A) Homogeneous ODE of degree 2
 - (B) Homogeneous ODE of degree 0
 - (C) Homogeneous ODE of degree 1
 - (D) Non homogeneous ODE

20. $\frac{dy}{dx} = \frac{(x+y)+1}{2(x+y)+1}$ reduces to variable separable form if :
- (A) $x - y = v$
 - (B) $x + y = v$
 - (C) $x^2 + y^2 = v$
 - (D) $xy = v$
21. Integrating factor in ODE $\frac{dy}{dx} + ay = e^{mx}$ is :
- (A) e^{mx}
 - (B) e^x
 - (C) e^a
 - (D) e^{ax}
22. Particular integral for $F(D)y = f(x)$ is :
- (A) $\frac{1}{F(D)} f(x)$
 - (B) $\frac{1}{F(D)} f(x) + c$ ($c \neq 0$)
 - (C) $\frac{1}{F(D)} f(x) + 1$
 - (D) NOT
23. Solution of $\frac{d^2y}{dx^2} - y = 0$ is :
- (A) $y = c_1 e^x + c_2 e^{-x}$
 - (B) $y = c_1 e^x + c_2 e^x$
 - (C) $y = c_1 e^x$
 - (D) $(c_1 + c_2 x) e^x$
24. Maximum value of directional derivative of $\phi = 2x^2 + 3y^2 + 5z^2$ at $(1, 1, -1)$ is :
- (A) 10
 - (B) -4
 - (C) $\sqrt{152}$
 - (D) 152

25. Divergence of $\vec{r}$ is :
- (A) 1
 - (B) 0
 - (C) 3
 - (D) 4
26. $\sin^{-1} x + \cos^{-1} x$ is :
- (A) $\frac{\pi}{2}$
 - (B) $\frac{-\pi}{3}$
 - (C) $\frac{\pi}{3}$
 - (D) NOT
27. Solution of $x dx - y dy = 0$ is :
- (A) $x = y + c$
 - (B) $x^2 - y^2 = c$
 - (C) $x - y = c$
 - (D) NOT
28. $\frac{dy}{dx} + 2y = x^2$ is :
- (A) Linear ODE in x
 - (B) Linear in y
 - (C) Linear in both variable x and y
 - (D) NOT
29. Degree of homogeneous ODE $(x^2 - y^2)dx + 3xy dy = 0$ is :
- (A) 1
 - (B) 2
 - (C) 0
 - (D) NOT

30. ODE $y \frac{dy}{dx} = x$ is :
- (A) Linear in y
 - (B) Non Linear in y
 - (C) Order of ODE is 2
 - (D) Degree of ODE is 2
31. Order of ODE $\left(\frac{dy}{dx}\right)^2 + y = 0$ is :
- (A) 1
 - (B) 2
 - (C) Not defined
 - (D) NOT
32. Integrating factor in linear ODE $\frac{dy}{dx} + Py = Q$ is :
- (A) $\int_e P dx$
 - (B) $\int_e P dy$
 - (C) $\int_e P dx + c$
 - (D) $\int_e P dy + c$
33. Degree of ODE $\left(\frac{d^2y}{dx^2}\right)^2 + \frac{d^2y}{dx^2} + y = 0$ is :
- (A) 1
 - (B) 2
 - (C) 4
 - (D) NOT
34. $f(x, y) = x^2 + y^2 + xy$ is homogeneous function of degree :
- (A) 1
 - (B) 0
 - (C) 2
 - (D) NOT

35. $\frac{dy}{dx} + Py = Qy^n$ ($n \neq 0, 1$) is :
- (A) Linear in y
 - (B) Non Linear in x
 - (C) Bernoulli equation
 - (D) NOT
36. ODE $M dx + N dy = 0$ is exact if :
- (A) $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$
 - (B) $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$
 - (C) $\frac{\partial M}{\partial x} = \frac{\partial N}{\partial y}$
 - (D) $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial y}$
37. $\frac{dy}{dx} + Py = Qy^n$ is linear ODE if :
- (A) $n = 1$
 - (B) $n = 2$
 - (C) $n = 0$
 - (D) $n = -1$
38. In Fourier series of $f(x)$, $f(x)$ is :
- (A) Periodic
 - (B) Non periodic
 - (C) Discontinuous
 - (D) Non differentiable
39. In Fourier series, a_0, a_n, b_n are obtained by :
- (A) Euler's formula
 - (B) Laplace formula
 - (C) Taylor's formula
 - (D) Maclaurin's formula

40. If $f(x)$ is periodic with period $\omega = 2l$ then Fourier coefficient :

(A) $a_0 = \frac{1}{4l} \int_a^b f(x) dx$

(B) $a_0 = \frac{1}{l} \int_a^b f(x) dx$

(C) $a_0 = \frac{4}{l} \int_a^b f(x) dx$

(D) NOT

41. Argument value of $z = -1 - i$ is :

(A) $\frac{\pi}{4}$

(B) $\frac{-\pi}{4}$

(C) $\frac{-3\pi}{4}$

(D) $\frac{3\pi}{4}$

42. Modulus value of $z = a + ib$ is :

(A) $a^2 + b^2$

(B) $\sqrt{a^2 + b^2}$

(C) $\sqrt{a^2 - b^2}$

(D) $a^2 - b^2$

43. Sequence $\{3, 3, 3, 3, \dots\}$ is :

(A) Bounded above only

(B) Bounded below only

(C) Bounded above and bounded below

(D) Unbounded

44. If sequence $\langle a_n \rangle$ is convergent then $\lim_{n \rightarrow \infty} a_n$ is :

(A) unique

(B) Not unique

(C) Always more than one limits

(D) NOT

45. n^{th} partial sum in the series $\sum_{n=1}^{\infty} a_n$ is :
- (A) $a_1 + a_2 + a_3 + \dots + a_{n-1}$
 (B) $a_1 + a_2 + a_3 + \dots + a_n$
 (C) $a_1 + a_2 + a_3 + \dots + a_{n+1}$
 (D) $a_0 + a_1 + a_2 + \dots + a_n$
46. Sequence $\{1, -1, 1, -1, 1, -1, \dots\}$ is
- (A) Convergent
 (B) Divergent
 (C) Oscillatory
 (D) Not convergent
47. Series $\sum_{n=0}^{\infty} x^n$ is oscillatory if :
- (A) $x = 1$
 (B) $x = -1$
 (C) $x = 2$
 (D) $x = 0$
48. Series $\sum_{n=1}^{\infty} (-1)^n$ is :
- (A) Convergent
 (B) Divergent
 (C) Oscillatory
 (D) NOT
49. Series $\sum_{n=1}^{\infty} \left(\frac{1}{n}\right)^{2022}$ is :
- (A) Convergent
 (B) Divergent
 (C) Oscillatory
 (D) Oscillatory finitely
50. If $z = -1 + i$ then modulus amplitude form of z is :
- (A) $z = \sqrt{2} e^{(5\pi/4)i}$
 (B) $z = \sqrt{2} e^{(-5\pi/4)i}$
 (C) $z = \sqrt{2} e^{(\pi/4)i}$
 (D) $z = \sqrt{2} e^{(-\pi/4)i}$

51. Roots of equation $z^2 + 1 = 0$ are :
- (A) $i, -i$
 - (B) $1, i$
 - (C) $1, -1$
 - (D) $-1, i$
52. Real part of $f(z) = 4 + i$ is :
- (A) 1
 - (B) 4
 - (C) $4 + i$
 - (D) i
53. $\overline{z_1 + z_2}$ is equal to :
- (A) $\overline{z_1} - \overline{z_2}$
 - (B) $\overline{z_1} + \overline{z_2}$
 - (C) $z_1 - z_2$
 - (D) NOT
54. Value of $\sqrt{-100}$ is :
- (A) 10
 - (B) -10
 - (C) $10i$
 - (D) $-10i$
55. In Fourier series of $f(x)$, coefficients a_0, a_n, b_n are called :
- (A) Laplace coefficients
 - (B) Fourier coefficients
 - (C) Taylor's coefficients
 - (D) Not
56. If $f(x) = \sin x$ is given in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, then value of a_n in Fourier series :
- (A) 0
 - (B) 1
 - (C) 2
 - (D) 3

57. Differential equation $\frac{3dy}{dx} + \frac{2}{x+1}y = \frac{x^3}{y^2}$ is reduced to linear for transformation :
- (A) $y = x^3$
 - (B) $y^2 = t$
 - (C) $y^3 = t$
 - (D) NOT
58. $\int_c^{c+2\pi} \sin nx \, dx$ is :
- (A) π
 - (B) c
 - (C) 0
 - (D) 2π
59. If $f(x) = \cos x$ in $[-l, l]$ then Fourier coefficient b_n is :
- (A) 1
 - (B) l
 - (C) $-l$
 - (D) 0
60. Fourier coefficients a_0, a_n, b_n are obtained by :
- (A) Taylor's Formula
 - (B) Maclaurin's Formula
 - (C) Euler's Formula
 - (D) Laplace Formula
61. $\frac{1}{D^2} \sin 2x$ is equal to :
- (A) $\frac{-1}{2} \sin 2x$
 - (B) $\frac{-1}{16} \sin 2x$
 - (C) $\frac{1}{4} \sin 2x$
 - (D) $\frac{-1}{4} \sin 2x$

62. $\frac{1}{D+1}e^x$ is :
- (A) e^x
 - (B) $\frac{1}{2}e^x$
 - (C) $\frac{1}{4}e^x$
 - (D) $\frac{-1}{4}e^x$
63. If $f(x) = x$ is periodic in $[0, 2\pi]$ then Fourier coefficient a_0 is :
- (A) $a_0 = \pi$
 - (B) $a_0 = 2\pi$
 - (C) $a_0 = \pi/2$
 - (D) NOT
64. Modulus value of $z = x + iy$ is :
- (A) $\sqrt{x^2 + y}$
 - (B) $\sqrt{x^2 + y^2}$
 - (C) $\tan^{-1}\left(\frac{y}{x}\right)$
 - (D) $\sqrt{x^2 - y^2}$
65. Cube roots of unity are :
- (A) $1, w, w^2$
 - (B) $1, -1, 0$
 - (C) $1, 1, 1$
 - (D) Not
66. Argument value of $z = 1 + i$ is :
- (A) $\frac{\pi}{2}$
 - (B) $\frac{\pi}{4}$
 - (C) 0
 - (D) $\frac{-\pi}{4}$

67. If $(x + 5) + i(y + 3) = 0$ then :
- (A) $x = 5, y = 3$
 - (B) $x = -5, y = 3$
 - (C) $x = 5, y = -3$
 - (D) $x = -5, y = -3$
68. Differential equation $(1 + 4xy + 2y^2)dx + (1 + 4xy + 2x^2)dy = 0$ is :
- (A) Linear in x
 - (B) Linear in y
 - (C) Exact ODE
 - (D) Homogeneous ODE
69. Order of ODE $\left(\frac{dy}{dx}\right)^2 + y = 0$ is :
- (A) 1
 - (B) 2
 - (C) Not defined
 - (D) -1
70. Degree of ODE $\sin\left(\frac{dy}{dx}\right) = 1$ is :
- (A) 1
 - (B) 0
 - (C) 2
 - (D) Does not exist
71. Maximum rate of increase of scalar point function $\phi(x, y, z)$ is :
- (A) $\nabla\phi$
 - (B) $-|\nabla\phi|$
 - (C) $|\nabla\phi|$
 - (D) $-\nabla\phi$

72. If $\phi(x, y, z) = x + y$, then directional derivative of $\phi(x, y, z)$ in the direction of $\vec{a} = i$ is :
- (A) 0
(B) 1
(C) -1
(D) i
73. If $\vec{v} = \vec{w} \times \vec{r}$ then $\text{curl } \vec{v}$ is :
- (A) $\vec{w}$
(B) $\vec{w} \times \vec{r}$
(C) $2\vec{w}$
(D) NOT
74. If $\vec{r} = xi + yj + zk$ then $\text{curl } \vec{r}$ is equal to :
- (A) 1
(B) $\vec{r}$
(C) 0
(D) 1
75. If $\phi(x, y, z) = x + y + z$ then gradient of ϕ is :
- (A) $i + j + k$
(B) $i + j$
(C) $x + y + z$
(D) 0
76. Which of the following is true ?
- (A) $|z_1 + z_2| \geq |z_1| + |z_2|$
(B) $|z_1 + z_2| = |z_1| + |z_2|$
(C) $|z_1 + z_2| \leq |z_1| + |z_2|$
(D) NOT

77. If $(2x + 2) + i(y - 5) = 0$ then :
- (A) $x = 1, y = 5$
 - (B) $x = -1, y = -5$
 - (C) $x = 1, y = -5$
 - (D) $x = -1, y = 5$
78. If $f(x) = \sin x \cos 2x$ then $f(x)$ is :
- (A) Even function
 - (B) Odd function
 - (C) Neither even nor odd
 - (D) NOT
79. Infinite series $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{c}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{c}\right)$ is :
- (A) Fourier series
 - (B) Laplace series
 - (C) Taylor's series
 - (D) Not
80. If $f(x) = \cos x$ is defined in $[-1, 1]$ then Fourier coefficient b_n is :
- (A) 4
 - (B) -1
 - (C) +1
 - (D) 0
81. Real part of $f(z) = \sin z$ is :
- (A) $\sin x \cos y$
 - (B) $\sin x \cos hy$
 - (C) $\sin hx \cos y$
 - (D) $\sin x \sin hy$
82. Euler's Formula is :
- (A) $e^{i\theta} = \cos \theta + i \sin \theta$
 - (B) $e^{-i\theta} = \cos \theta + i \sin \theta$
 - (C) $e^{i\theta} = 2 \cos \theta + \sin \theta$
 - (D) $e^{-i\theta} = \cos \theta - \sin \theta$

83. A vector which is perpendicular to both vectors $\vec{a}$ and $\vec{b}$ is :

(A) $|\vec{a} \times \vec{b}|$

(B) $\vec{a} \times \vec{b}$

(C) $|2\vec{a} \times \vec{b}|$

(D) $|\vec{b} \times \vec{a}|$

84. $\nabla f(r)$ is :

(A) $f'(r)$

(B) $f'(r)\nabla r$

(C) $f'(r)\vec{r}$

(D) $\frac{f'(r)}{r}$

85. Gradient of $|\vec{r}|$ is :

(A) $\frac{\vec{r}}{r}$

(B) $\vec{r}$

(C) $\frac{\vec{r}}{r^2}$

(D) r

86. For solenoidal field vector $\vec{v}$, divergence of $\vec{v}$ is :

(A) Non zero

(B) Zero

(C) Never zero

(D) NOT

87. $\frac{d}{dt}(\vec{a} \times \vec{b}) =$

(A) $\vec{a} \times \frac{d\vec{b}}{dt} + \frac{d\vec{a}}{dt} \times \vec{b}$

(B) $\vec{a} \times \frac{d\vec{b}}{dt} + \frac{d\vec{a}}{dt} \vec{b}$

(C) $\vec{a} \times \frac{d\vec{b}}{dt} + \vec{b} \times \frac{d\vec{a}}{dt}$

(D) NOT

88. Curl of a vector field is :
- (A) Scalar
 - (B) Vector
 - (C) Never be a vector
 - (D) NOT
89. $\tan^{-1}\left(\frac{4}{7}\right) + \tan^{-1}\left(\frac{3}{11}\right)$ is :
- (A) $\frac{\pi}{4}$
 - (B) $\frac{-\pi}{4}$
 - (C) 1
 - (D) NOT
90. Imaginary part of $\frac{3-2i}{7+4i}$ is :
- (A) $\frac{2}{5}$
 - (B) $\frac{-2}{5}$
 - (C) $\frac{1}{5}$
 - (D) NOT
91. Locus of $|z| < 1$ is :
- (A) $x^2 + y^2 = 1$
 - (B) $x^2 + y^2 > 1$
 - (C) $x^2 + y^2 < 1$
 - (D) $x + y < 1$
92. $\frac{e^{iz} - e^{-iz}}{2i}$ is equal to :
- (A) $\cos z$
 - (B) $\cos hz$
 - (C) $\sin z$
 - (D) $\sin hz$

93. If $f(x)$ is periodic function with period $2c$ then Fourier coefficient a_0 is :

(A) $\frac{1}{4c} \int_a^b f(x) dx$

(B) $\frac{1}{3c} \int_a^b f(x) dx$

(C) $\frac{1}{c} \int_a^b f(x) dx$

(D) NOT

94. ODE $\frac{dy}{dx} = \frac{x^2+y^2}{xy}$ is :

(A) Homogeneous ODE

(B) Non homogeneous ODE

(C) Linear in x

(D) Linear in y

95. Sequence $\langle \frac{1}{n} \rangle$ is :

(A) Bounded above

(B) Bounded below

(C) Neither bounded above nor bounded below

(D) Option (A) and (B) both

96. Sequence $\langle a_n \rangle = \langle n \rangle$ has :

(A) Infinitely many terms

(B) Finitely many terms

(C) n terms

(D) NOT

97. Sequence $\langle \frac{1}{n+1} \rangle$ converges to :
- (A) 1
 - (B) 0
 - (C) ∞
 - (D) -1
98. Sequence $\langle a_n \rangle = \{5, 5, 5, \dots\}$ i.e. $a_n = 5 \ \forall \ n \in N$ is :
- (A) Constant sequence
 - (B) Not bounded above
 - (C) Not bounded below
 - (D) Not bounded
99. Complementary function for ODE $(D^2 - 1)y = 0$, where $D \equiv \frac{d}{dx}$ is :
- (A) $(c_1 + c_2 x)e^x$
 - (B) $c_1 e^x + c_2 e^{-x}$
 - (C) e^x
 - (D) e^{-x}
100. Complementary function for $(D^2 + 1)y = 0$ is :
- (A) $c_1 \cos x$
 - (B) $c_2 \sin x$
 - (C) $c_1 + c_2 \cos x$
 - (D) $c_1 \cos x + c_2 \sin x$

Rough Work / रफ कार्य

DO NOT OPEN THE QUESTION BOOKLET UNTIL ASKED TO DO SO

1. Examinee should enter his / her roll number, subject and Question Booklet Series correctly in the O.M.R. sheet, the examinee will be responsible for the error he / she has made.
 2. **This Question Booklet contains 100 questions, out of which only 75 Question are to be Answered by the examinee. Every question has 4 options and only one of them is correct. The answer which seems correct to you, darken that option number in your Answer Booklet (O.M.R ANSWER SHEET) completely with black or blue ball point pen. If any examinee will mark more than one answer of a particular question, then the answer will be marked as wrong.**
 3. Every question has same marks. Every question you attempt correctly, marks will be given according to that.
 4. Every answer should be marked only on Answer Booklet (O.M.R ANSWER SHEET). Answer marked anywhere else other than the determined place will not be considered valid.
 5. Please read all the instructions carefully before attempting anything on Answer Booklet **(O.M.R ANSWER SHEET)**.
 6. After completion of examination, please hand over the **O.M.R. SHEET** to the Examiner before leaving the examination room.
 7. There is no negative marking.
- Note:** On opening the question booklet, first check that all the pages of the question booklet are printed properly in case there is an issue please ask the examiner to change the booklet of same series and get another one.